

## ANTG SOLACE

### Consumer Medicine Information (CMI) summary

The full CMI on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

#### 1. WHY AM I USING ANTG SOLACE?

ANTG Solace contains active ingredients from whole dried *Cannabis Sativa* flower. ANTG Solace is not approved for any specific indication and should only be used to treat conditions or symptoms advised by your doctor.

Your doctor might recommend its use to reduce nausea, vomiting, chronic non cancer pain, cancer related pain, spasticity from neurological conditions and muscle spasms, improve sleep and appetite, depression and anxiety. Your doctor might prescribe ANTG Solace for another purpose.

For more information, see [Section 1. Why am I using ANTG Solace?](#) in the full CMI.

#### 2. WHAT SHOULD I KNOW BEFORE I USE ANTG SOLACE?

Do not use if you have ever had an allergic reaction to cannabis or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see [Section 2. What should I know before I use ANTG Solace?](#) in the full CMI.

#### 3. WHAT IF I AM TAKING OTHER MEDICINES?

Some medicines may interfere with ANTG Solace and affect how it works. ANTG Solace may affect how other medicines work.

A list of these medicines is in [Section 3. What if I am taking other medicines?](#) in the full CMI.

#### 4. HOW DO I USE ANTG SOLACE?

Use the amount of ANTG Solace directed by your doctor. Follow your doctor's instructions and use ANTG Solace until your doctor tells you to stop.

More instructions can be found in [Section 4. How do I use ANTG Solace?](#) in the full CMI.

#### 5. WHAT SHOULD I KNOW WHILE USING ANTG SOLACE?

<b>Things you should do</b>	<ul style="list-style-type: none"> <li>Remind any doctor, dentist or pharmacist you visit that you are using ANTG Solace.</li> <li>You should use ANTG Solace under medical supervision.</li> <li>Talk to your healthcare provider if you do not feel better, or if you feel worse.</li> </ul>
<b>Things you should not do</b>	<ul style="list-style-type: none"> <li>Do not change your dosage without first checking with your doctor.</li> </ul>
<b>Driving or using machines</b>	<ul style="list-style-type: none"> <li>Do not drive or operate heavy or dangerous machines while being treated on this product.</li> </ul>
<b>Drinking alcohol</b>	<ul style="list-style-type: none"> <li>Alcohol may increase the effects of ANTG Solace and you should avoid taking it while taking this medicine. Alcohol taken in combination with ANTG Solace can impair your cognitive abilities, coordination, reaction time and motor skills.</li> </ul>
<b>Looking after your medicine</b>	<ul style="list-style-type: none"> <li>Store below 25°C and protect from light.</li> </ul>

For more information, see [Section 5. What should I know while using ANTG Solace?](#) in the full CMI.

#### 6. ARE THERE ANY SIDE EFFECTS?

There are a number of side effects associated with this medicine. It is important to be aware of them so that you can identify any symptoms if they occur (see the full CMI for more details). Common, but usually mild side effects known from medicinal cannabis treatment (both cannabidiol (CBD) and delta-9-tetrahydrocannabinol (THC)) include fatigue and sedation, dizziness, nausea, dry mouth, and increase or decrease in appetite. Serious side effects of THC (and products high in THC) include convulsions, feeling 'high', depression, confusion, hallucinations, paranoid delusions, psychosis (severe mental condition where a person loses contact with reality, and is unable to think and judge clearly), cognitive distortion (having thoughts that are not true), and allergy symptoms.

For more information, including what to do if you have any side effects, see [Section 6. Are there any side effects?](#) in the full CMI.

## ANTG SOLACE

### (/i:v/) Active ingredient:

Total THC [THC+THCA (as THC)] 200mg/g (20% w/w)

Total CBD [CBD+CBDA (as CBD)] <10mg/g (<1% w/w)

### Consumer Medicine Information (CMI)

This leaflet provides important information about using ANTG Solace. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ANTG Solace.**

#### WHERE TO FIND INFORMATION IN THIS LEAFLET:

- [1. Why am I using ANTG Solace?](#)
- [2. What should I know before I use ANTG Solace?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use ANTG Solace?](#)
- [5. What should I know while using ANTG Solace?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

#### 1. WHY AM I USING ANTG SOLACE?

ANTG Solace contains the active ingredients from whole dried *Cannabis Sativa* flower cured and trimmed from a hybrid cultivar of the *Cannabis Sativa* plant. The trichomes (glands) on the cannabis sativa flower are abundant with phytocannabinoid compounds. When taken internally, phytocannabinoids interact with the body's receptors to produce psychoactive effects (changes to a person's mental state, such as behavior, mood, thoughts or perception) and therapeutic effects. ANTG Solace is not approved for any specific indication and should only be used to treat conditions or symptoms advised by your doctor.

Your doctor might recommend its use to reduce nausea, vomiting, chronic non cancer pain, cancer related pain, spasticity from neurological conditions and muscle spasms, improve sleep and appetite, and manage seizures, depression and anxiety. Your doctor might prescribe ANTG Solace for another purpose. This medicine is only available with a doctor's prescription. Possession without authority is illegal.

#### 2. WHAT SHOULD I KNOW BEFORE I USE ANTG SOLACE?

##### WARNINGS

##### DO NOT USE ANTG SOLACE IF:

- You are allergic to any cannabis product or any of the ingredients listed at the end of this leaflet (Always check the ingredients to make sure you can use this medicine)
- You are less than 18 years old
- You have serious liver or kidney problems
- You have pre-existing heart problems, such as narrowing of the arteries (ischaemic or coronary heart disease), irregular heartbeat, poorly controlled high blood pressure, or have had previous heart attack or heart failure

#### CHECK WITH YOUR DOCTOR IF YOU:

- Have or have had any other medical conditions, especially the following:
  - Kidney problems
  - Liver problems
  - Serious mental disorders, such as schizophrenia or another psychotic disorder
  - Panic attacks
  - Lung problems
  - Heart problems
  - Epilepsy or recurrent seizures
- You have a history of dependence or addictive behaviour associated with prescription or non-prescription products
- You are 65 years or older

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under [Section 6. Are there any side effects?](#)

#### PREGNANCY AND BREASTFEEDING

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed. Cannabinoids can pass into breastmilk and may affect foetal development.

Do not take this product whilst breastfeeding until you and your doctor have discussed the risks and benefits involved.

#### USE IN CHILDREN

ANTG Solace is not recommended for use in children or adolescents under 18 years of age.

#### MEDICAL SUPERVISION OF TREATMENT WITH ANTG SOLACE

ANTG Solace treatment should be initiated and supervised by a qualified healthcare provider experienced in the treatment of the symptom or disorder for which it is prescribed.

#### 3. WHAT IF I AM TAKING OTHER MEDICINES?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Some medicines may interfere with ANTG Solace and affect how it works. ANTG Solace may affect how other medicines work. Tell your doctor if you are using medicines for:**

- Cold or allergy (medicines that make you drowsy, such as antihistamines)
- Pain (medicines known as opioids, such as oxycodone, hydrocodone, codeine, and morphine)
- Muscle-related conditions, such as muscle spasms, overactive bladder, movement disorders (Parkinson's disease), and asthma (medicines known as anticholinergics, e.g. tolterodine, oxybutynin, darifenacin, solifenacin, benzotropine mesilate, trihexyphenidyl, ipratropium bromide, tiotropium bromide)
- Sleeping problems (sleeping pills or sedatives/hypnotics, e.g. benzodiazepine, zopiclone, zolpidem)
- Epilepsy, seizures, or nerve pain (e.g. carbamazepine, phenytoin, and phenobarbital)
- Attention Deficit Hyperactivity Disorder (ADHD) or narcolepsy (medicines such as amphetamines that may increase your heart rate and blood pressure)

- Bacterial or fungal infections (e.g. rifampicin, clarithromycin, ketoconazole, itraconazole, amphotericin B)
- Suppression of the immune system (e.g. cyclosporine)
- Thinning of your blood (medicines known as coumarins, e.g. warfarin)

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ANTG Solace.**

#### 4. HOW DO I USE ANTG SOLACE?

##### HOW MUCH TO USE

- Use the amount of ANTG Solace directed by your doctor.
- The amount of ANTG Solace your doctor tells you to use each day will depend on your condition, whether you are taking any other medicines, as well as how you respond to and tolerate the product.
- Your doctor may recommend that you start with a low dose of ANTG Solace and then slowly increase the dose to the lowest amount needed to improve your symptoms.
- Follow the instructions provided and use ANTG Solace until your doctor tells you to stop.

##### WHEN TO TAKE ANTG SOLACE

- ANTG Solace can be taken with or without food.
- Taking ANTG Solace under similar conditions (e.g. before or after food/meals) each day may help achieve a more consistent effect.

##### HOW TO TAKE ANTG SOLACE

- This product should be administered as directed by your doctor.
- Effects from ANTG Solace may vary depending on the dose taken, how long you have been taking this medicine, the amount and type of food you have had when taking this medicine, and your individual response.

##### Tea Preparation for Oral Consumption

1. Bring 500 ml (0.5 litres) of water to the boil in a covered vessel
2. Granulate product (using a grinder is easiest) into fine pieces
3. Add 0.5 grams to vessel
4. Turn down the heat and simmer the tea for 15 minutes with the lid on the vessel
5. Remove the covered vessel from the heat and strain the tea
6. Store the leftover tea in a thermos if you are to drink more of it on the same day

After preparation - as described above - add a sachet of coffee creamer powder to the warm tea. This prevents the active ingredients in the tea from sticking to the inside of the teapot or mug after cooling, which could reduce the efficacy of the tea.

##### IF YOU FORGET TO USE ANTG SOLACE

ANTG Solace should be used regularly at the same time each day. If you miss your dose at the usual time, take it as soon as you remember, then go back to taking this product as you would normally.

**If you have missed a dose by more than 4 hours, skip the dose you missed and take your next dose when you are meant to.**

**Do not take a double dose to make up for the dose you missed.**

##### IF YOU USE TOO MUCH ANTG SOLACE

If you think that you have used too much ANTG Solace, you may need urgent medical attention.

##### You should immediately:

- phone the New Zealand National Poisons Centre (**by calling 0800 764 766**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

#### 5. WHAT SHOULD I KNOW WHILE USING ANTG SOLACE?

##### THINGS YOU SHOULD DO

- You should use ANTG Solace under medical supervision to help determine if the product is working or if you are having any side effects from the treatment.
- Talk to your healthcare provider if you do not feel better, or if you feel worse

Remind any doctor, dentist or pharmacist you visit that you are using ANTG Solace.

##### THINGS YOU SHOULD NOT DO

- Do not change your dosage without first checking with your doctor.

##### TRAVELLING WITH ANTG SOLACE

- Before travelling to another country (this includes your final destination and any in transit countries), consult with your doctor, country consulates, The Ministry of Foreign Affairs and Trade, insurance company and/or pharmacist to confirm if it is legal for you to travel with this product.

##### DRIVING OR USING MACHINES

**Do not drive or operate heavy or dangerous machines while being treated on this product. Talk to your doctor on when to resume these activities after completing treatment.**

##### DRINKING ALCOHOL

**Tell your doctor if you drink alcohol.**

Alcohol may increase the effects of ANTG Solace and you should avoid taking it while taking this medicine. Alcohol taken in combination with ANTG Solace can impair your cognitive abilities, coordination, reaction time and motor skills.

##### LOOKING AFTER YOUR MEDICINE

- Store below 25°C.
- Follow the instructions in the bottle on how to take care of your medicine properly.
- Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:
  - in the bathroom or near a sink, or
  - in the car or on window sills.

**Keep it where young children cannot reach it.**

##### GETTING RID OF ANY UNWANTED MEDICINE

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

#### 6. ARE THERE ANY SIDE EFFECTS?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

#### SIDE EFFECTS

Side effects	What to do
<b>Changes to mental status:</b> <ul style="list-style-type: none"> <li>• Poor short-term memory</li> <li>• Lack of concentration</li> <li>• Confusion</li> <li>• Abnormal thoughts</li> <li>• Poor coordination and balance</li> <li>• Dizziness</li> </ul>	Speak to your doctor if you have any of these less serious side effects and they worry you.
<b>Personality or behavioural changes:</b> <ul style="list-style-type: none"> <li>• Anxiety</li> <li>• Loss of inhibition</li> <li>• Quiet and reflective mood, or other unusual changes in mood</li> <li>• Changes in appetite</li> <li>• Spontaneous laughter</li> <li>• Sleepiness/feeling tired</li> </ul>	
<b>Eye-related:</b> <ul style="list-style-type: none"> <li>• Bloodshot eyes</li> <li>• Altered visual perception</li> </ul>	
<b>Ear-related:</b> <ul style="list-style-type: none"> <li>• Altered hearing perception</li> </ul>	
<b>General body-related:</b> <ul style="list-style-type: none"> <li>• Phlegm and cough</li> <li>• Dry mouth</li> <li>• Malaise (feeling unwell) or a general feeling of discomfort</li> <li>• Nausea</li> <li>• Muscle weakness</li> </ul>	Speak to your doctor if you have any of these less serious side effects and they worry you.

#### SERIOUS SIDE EFFECTS

Serious side effects	What to do
<b>Changes to mental status or personality:</b> <ul style="list-style-type: none"> <li>• Depression or thoughts of suicide</li> <li>• Paranoia</li> <li>• Seeing or hearing things that are not there</li> <li>• Irrational thinking</li> <li>• Hallucinations</li> <li>• Addiction/dependence</li> </ul>	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.
<b>Heart-related:</b> <ul style="list-style-type: none"> <li>• Rapid or irregular heartbeat</li> <li>• Signs of low blood pressure, such as light-headedness, dizziness or fainting</li> </ul>	
<b>Nervous system-related:</b> <ul style="list-style-type: none"> <li>• Seizures or convulsions (if you do not have epilepsy)</li> </ul>	
<b>Allergic reaction-related:</b> <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Wheezing or difficulty breathing</li> <li>• Swelling of the face, lips, tongue or other parts of the body</li> <li>• Rash, hives or swelling on the skin</li> </ul>	

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

#### REPORTING SIDE EFFECTS

After you have received medical advice for any side effects you experience, you can report side effects to your doctor, pharmacist, NUBU Pharmaceuticals and/or the New Zealand Pharmacovigilance Centre by visiting [www.otago.ac.nz/carm](http://www.otago.ac.nz/carm).

By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

#### 7. PRODUCT DETAILS

This medicine is only available with a doctor's prescription.

##### WHAT ANTG SOLACE CONTAINS

<b>Active ingredient (main ingredient)</b>	Each bottle has 10 g whole dried <i>Cannabis Sativa</i> flower, which contains 2000mg of delta-9- tetrahydrocannabinol (20% THC), <100mg cannabidiol (<1% CBD) and other minor cannabinoids.
<b>Other ingredients (inactive ingredients)</b>	N/A

Do not take this medicine if you are allergic to any of these ingredients.

##### WHAT ANTG SOLACE LOOKS LIKE

Approximately 1–5 cm, brown-green clustered flowers with a characteristic cannabis smell. Available in a white High Density Polyethylene jar with a child-resistant cap and tamper proof seal.

##### WHO DISTRIBUTES ANTG SOLACE

NUBU Pharmaceuticals  
Auckland, New Zealand.

Email: [info@nubupharma.com](mailto:info@nubupharma.com)  
Ph: 0800 46 33 22

Date of Preparation March 2022. NUBU025V2.2